

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PIK3R2 / Phosphoinositide-3-kinase regulatory subunit 2 Antibody, IgG2b, Clone: [PIK3R2/292], Mouse, Monoclonal NBT-5296-MSM2-P0

| | |
|--------------------------|---|
| Artikelname | PIK3R2 / Phosphoinositide-3-kinase regulatory subunit 2 Antibody, IgG2b, Clone: [PIK3R2/292], Mouse, Monoclonal |
| Artikelnummer | NBT-5296-MSM2-P0 |
| Hersteller Artikelnummer | 5296-MSM2-P0 |
| Alternativnummer | NBT-5296-MSM2-P0-20,NBT-5296-MSM2-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human PIK3R2 protein (around aa 520-720) (exact sequence is proprietary) |
| Produktbeschreibung | Phosphatidylinositol 3-kinase (PI 3-kinase) is composed of p85 and p110 subunits. p85 lacks PI 3-kinase activity and acts as an adapter, coupling p110 to activated protein tyrosine kinase. Two forms of p85 have been described (p85 and p85), each poss... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [PIK3R2/292] |
| Molekulargewicht | 85kDa |

| | |
|------------------------|---|
| Isotyp | IgG2b |
| NCBI | 5296 |
| UniProt | O00459 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |