

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Perforin-1 (Pore Forming Protein) (Apoptosis Marker) Antibody, Clone: [PRF1/8662R], Rabbit, Monoclonal NBT-5551-RBM9-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant Perforin-1 (Pore Forming Protein) (Apoptosis Marker) Antibody, Clone: [PRF1/8662R], Rabbit, Monoclonal |
| Artikelnummer | NBT-5551-RBM9-P0 |
| Hersteller Artikelnummer | 5551-RBM9-P0 |
| Alternativnummer | NBT-5551-RBM9-P0-20,NBT-5551-RBM9-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Perforin-1 protein fragment (around aa 413-552) (exact sequence is proprietary) |
| Produktbeschreibung | Perforin is a pore-forming protein that leads to osmotic lysis of the target cells and subsequently enables granzymes to enter the target cells and activate apoptosis. Perforin has structural and functional similarities to complement component 9 (C9)... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [PRF1/8662R] |
| Molekulargewicht | 75kDa |

| | |
|------------------------|--|
| NCBI | 5551 |
| UniProt | P14222 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |