

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Rb1 (Tumor Suppressor Protein) Antibody, Clone: [RB1/7080R], Rabbit, Monoclonal NBT-5925-RBM4-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant Rb1 (Tumor Suppressor Protein) Antibody, Clone: [RB1/7080R], Rabbit, Monoclonal |
| Artikelnummer | NBT-5925-RBM4-P0 |
| Hersteller Artikelnummer | 5925-RBM4-P0 |
| Alternativnummer | NBT-5925-RBM4-P0-20,NBT-5925-RBM4-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length Rb1 protein |
| Produktbeschreibung | Recognizes a 105kDa phosphoprotein, identified as retinoblastoma (Rb) gene product. Its epitope is localized between aa 703-772. It shows no cross reaction with p107 or p130. It specifically stains the nuclei of BT-20 cells and primary human foreskin... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [RB1/7080R] |
| Molekulargewicht | 105kDa |
| NCBI | 5925 |

| | |
|------------------------|--|
| UniProt | P06400 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |