

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant RBP4 / Retinol Binding Protein 4 Antibody, IgG1, Clone: [rRBP4/7372], Mouse, Monoclonal NBT-5950-MSM22-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant RBP4 / Retinol Binding Protein 4 Antibody, IgG1, Clone: [rRBP4/7372], Mouse, Monoclonal   |
| Artikelnummer            | NBT-5950-MSM22-P0   |
| Hersteller Artikelnummer | 5950-MSM22-P0   |
| Alternativnummer         | NBT-5950-MSM22-P0-20,NBT-5950-MSM22-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment (around aa29-148) of human RBP4 protein (exact sequence is proprietary)  |
| Produktbeschreibung      | Retinol (Vitamin A) is transported in the blood bound to its carrier protein, retinol-binding protein (RBP), also designated plasma retinol-binding protein (PRBP) or RBP4. A member of the lipocalin family, RBP conveys retinol from stores in the liver... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rRBP4/7372]  |
| Molekulargewicht         | 25kDa   |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| NCBI                   | <a href="#">5950</a>  |
| UniProt                | <a href="#">P02753</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |