

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody, IgG1, Clone: [100/D5+124], Mouse, Monoclonal NBT-596-MSM4-P0**

Artikelname	Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody, IgG1, Clone: [100/D5+124], Mouse, Monoclonal
Artikelnummer	NBT-596-MSM4-P0
Hersteller Artikelnummer	596-MSM4-P0
Alternativnummer	NBT-596-MSM4-P0-20,NBT-596-MSM4-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 (100/D5 & 124)
Produktbeschreibung	This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 lpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 lpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular ly...
Klonalität	Monoclonal
Klon-Bezeichnung	[100/D5+124]
Molekulargewicht	25-26kDa

Isotyp	IgG1
NCBI	<a href="#">596</a>
UniProt	<a href="#">P10415</a>
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris wit