

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Bcl-6 (Follicular Lymphoma Marker) Antibody, IgG1, Clone: [rBCL6/1718], Mouse, Monoclonal NBT-604-MSM14-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant Bcl-6 (Follicular Lymphoma Marker) Antibody, IgG1, Clone: [rBCL6/1718], Mouse, Monoclonal |
| Artikelnummer | NBT-604-MSM14-P1ABX |
| Hersteller Artikelnummer | 604-MSM14-P1ABX |
| Alternativnummer | NBT-604-MSM14-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity wherea... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [rBCL6/1718] |
| Molekulargewicht | 95kDa |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| NCBI | 604 |
| UniProt | P41182 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution |