

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant ROS1 Antibody, Clone: [ROS1/7999R], Rabbit, Monoclonal NBT-6098-RBM3-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant ROS1 Antibody, Clone: [ROS1/7999R], Rabbit, Monoclonal |
| Artikelnummer | NBT-6098-RBM3-P0 |
| Hersteller Artikelnummer | 6098-RBM3-P0 |
| Alternativnummer | NBT-6098-RBM3-P0-20,NBT-6098-RBM3-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment corresponding to cytoplasmic domain of human ROS1 protein |
| Produktbeschreibung | ROS1, an orphan receptor tyrosine kinase of the insulin receptor family, was initially identified as a homolog of v-ros from the UR2 sarcoma virus. ROS1 consists of a large extracellular domain that is composed of six fibronectin repeats, a transmemb... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [ROS1/7999R] |
| Molekulargewicht | 264kDa |
| NCBI | 6098 |

| | |
|------------------------|--|
| UniProt | P08922 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |