

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Monocyte Chemotactic Protein 2 (MCP2) / CCL8 Antibody, IgG1, Clone: [CCL8/3686], Mouse, Monoclonal NBT-6355-MSM6-P1ABX

| | |
|--------------------------|---|
| Artikelname | Monocyte Chemotactic Protein 2 (MCP2) / CCL8 Antibody, IgG1, Clone: [CCL8/3686], Mouse, Monoclonal |
| Artikelnummer | NBT-6355-MSM6-P1ABX |
| Hersteller Artikelnummer | 6355-MSM6-P1ABX |
| Alternativnummer | NBT-6355-MSM6-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human recombinant MCP2 / CCL8 protein fragment (around aa1-99) (exact sequence is proprietary) |
| Produktbeschreibung | This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfam... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CCL8/3686] |
| Molekulargewicht | 11kDa |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| NCBI | 6355 |
| UniProt | P80075 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |