

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Gamma-parvin (PARVG) Antibody, Clone: [PARVG/6312R], Rabbit, Monoclonal NBT-64098-RBM2-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant Gamma-parvin (PARVG) Antibody, Clone: [PARVG/6312R], Rabbit, Monoclonal |
| Artikelnummer | NBT-64098-RBM2-P0 |
| Hersteller Artikelnummer | 64098-RBM2-P0 |
| Alternativnummer | NBT-64098-RBM2-P0-20,NBT-64098-RBM2-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | GST-tagged recombinant protein corresponding to human Gamma-parvin |
| Produktbeschreibung | The parvin family, including -parvin, -parvin and -parvin, link integrins and associated proteins with intracellular pathways, which regulate actin cytoskeletal dynamics and cell survival. All three family members localize to focal adhesions and func... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [PARVG/6312R] |
| Molekulargewicht | 37kDa |

| | |
|------------------------|--|
| NCBI | 64098 |
| UniProt | Q9HBIO |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |