

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Fascin-1 (Reed-Sternberg CellMarker) Antibody, Clone: [FSCN1/6465R], Rabbit, Monoclonal NBT-6624-RBM5-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant Fascin-1 (Reed-Sternberg CellMarker) Antibody, Clone: [FSCN1/6465R], Rabbit, Monoclonal |
| Artikelnummer | NBT-6624-RBM5-P1ABX |
| Hersteller Artikelnummer | 6624-RBM5-P1ABX |
| Alternativnummer | NBT-6624-RBM5-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Full length recombinant human FSCN1 protein |
| Produktbeschreibung | Recognizes a protein of 55kDa, which is identified as fascin-1. Its actin binding ability is regulated by phosphorylation. Antibody to fascin-1 is a very sensitive marker for Reed-Sternberg cells and variants in nodular sclerosis, mixed cellularity, ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [FSCN1/6465R] |
| Molekulargewicht | 55kDa |
| NCBI | 6624 |

| | |
|------------------------|---|
| UniProt | Q16658 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |