

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD147 / EMMPRIN / Neurothelin Antibody, Clone: [BSG/6468R], Rabbit, Monoclonal NBT-682-RBM7-P1ABX

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Recombinant CD147 / EMMPRIN / Neurothelin Antibody, Clone: [BSG/6468R], Rabbit, Monoclonal |
| Artikelnummer | NBT-682-RBM7-P1ABX |
| Hersteller Artikelnummer | 682-RBM7-P1ABX |
| Alternativnummer | NBT-682-RBM7-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human BSG protein |
| Produktbeschreibung | This MAb recognizes extracellular epitope 2 within the N-terminal Ig domain of human CD147. It is expressed more intensely on thymocytes than on mature peripheral blood T cells. CD147 is important in spermatogenesis, embryo implantation, neural netwo... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BSG/6468R] |
| Molekulargewicht | 35kDa (non-reduced), 40kDa (reduced) |
| NCBI | 682 |

| | |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P35613 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Inhibits Homotypic Aggregation, Adhesion to Matrix Proteins, & Migration through Matrigel (Order w/o BSA / azide), Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (0.5-1ug/ml for 30 minutes at |