

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **pS2 / pNR-2 / TFF1 (Estrogen-Regulated Protein) Antibody, Clone: [TFF1/7891], Mouse, Monoclonal NBT-7031-MSM11-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | pS2 / pNR-2 / TFF1 (Estrogen-Regulated Protein) Antibody, Clone: [TFF1/7891], Mouse, Monoclonal   |
| Artikelnummer            | NBT-7031-MSM11-P0   |
| Hersteller Artikelnummer | 7031-MSM11-P0   |
| Alternativnummer         | NBT-7031-MSM11-P0-20,NBT-7031-MSM11-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment (around aa1-84) of human TFF1 protein (exact sequence is proprietary)  |
| Produktbeschreibung      | It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is located in the c-terminus of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gu... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [TFF1/7891]   |
| Molekulargewicht         | 6.5kDa  |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">7031</a>   |
| UniProt                | <a href="#">P04155</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |