

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **A20 / TNFAIP3 (Marker of Tamoxifen Resistance) Antibody, IgG2c, Clone: [TNFAIP3/2813], Mouse, Monoclonal NBT-7128-MSM3-P0**

Artikelname	A20 / TNFAIP3 (Marker of Tamoxifen Resistance) Antibody, IgG2c, Clone: [TNFAIP3/2813], Mouse, Monoclonal
Artikelnummer	NBT-7128-MSM3-P0
Hersteller Artikelnummer	7128-MSM3-P0
Alternativnummer	NBT-7128-MSM3-P0-20,NBT-7128-MSM3-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa 241-356) of human TNFAIP3 protein (exact sequence is proprietary)
Produktbeschreibung	A20 (also known as TNFAIP3) is a zinc finger protein and inhibits NF-kappa B activation as well as TNF-mediated apoptosis. The protein has both ubiquitin ligase and deubiquitinase activities, is involved in the cytokine-mediated immune and inflammato...
Klonalität	Monoclonal
Klon-Bezeichnung	[TNFAIP3/2813]
Molekulargewicht	~90kDa

Isotyp	IgG2c
NCBI	<a href="#">7128</a>
UniProt	<a href="#">P21580</a>
Formulierung	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) ,(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution