

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant TRPS1 (Transcriptional repressor GATA binding 1) Antibody, Clone: [TRPS1/8008R], Rabbit, Monoclonal NBT-7227-RBM8-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant TRPS1 (Transcriptional repressor GATA binding 1) Antibody, Clone: [TRPS1/8008R], Rabbit, Monoclonal   |
| Artikelnummer            | NBT-7227-RBM8-P1ABX   |
| Hersteller Artikelnummer | 7227-RBM8-P1ABX   |
| Alternativnummer         | NBT-7227-RBM8-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment (around aa900-1100) of human TRPS1 protein (exact sequence is proprietary)   |
| Produktbeschreibung      | Transcriptional repressor. Binds specifically to GATA sequences and represses expression of GATA-regulated genes at selected sites and stages in vertebrate development. Regulates chondrocyte proliferation and differentiation. Executes multiple functi... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [TRPS1/8008R]   |
| Molekulargewicht         | 142kDa  |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">7227</a>   |
| UniProt                | <a href="#">Q9UHF7</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Recombinant Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |