

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Carbonic Anhydrase VIII Antibody, IgG2a, Clone: [CA8/6572], Mouse, Monoclonal NBT-767-MSM2-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Carbonic Anhydrase VIII Antibody, IgG2a, Clone: [CA8/6572], Mouse, Monoclonal   |
| Artikelnummer            | NBT-767-MSM2-P1ABX  |
| Hersteller Artikelnummer | 767-MSM2-P1ABX  |
| Alternativnummer         | NBT-767-MSM2-P1ABX-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human CA8 protein   |
| Produktbeschreibung      | The protein encoded by this gene was initially named CA-related protein because of sequence similarity to other known carbonic anhydrase genes. However, the gene product lacks carbonic anhydrase activity (i.e., the reversible hydration of carbon diox... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [CA8/6572]  |
| Molekulargewicht         | 32.97kDa  |
| Isotyp                   | IgG2a   |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">767</a>   |
| UniProt                | <a href="#">P35219</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |