

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TIM3 / HAVCR2/ CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4025], Mouse, Monoclonal NBT-84868-MSM5-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | TIM3 / HAVCR2/ CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4025], Mouse, Monoclonal  |
| Artikelnummer            | NBT-84868-MSM5-P0   |
| Hersteller Artikelnummer | 84868-MSM5-P0   |
| Alternativnummer         | NBT-84868-MSM5-P0-20,NBT-84868-MSM5-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary)   |
| Produktbeschreibung      | TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cel tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 a... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [TIM3/4025]   |
| Molekulargewicht         | 30-60kDa  |

|                        |  |
|------------------------|--|
| Isotyp                 | IgG2b  |
| NCBI                   | <a href="#">84868</a>  |
| UniProt                | <a href="#">Q8TDQ0</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2 |