

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TIM3 / HAVCR2 / CD366 (Effector T-Cell Marker) Antibody, IgG2c, Clone: [TIM3/4028], Mouse, Monoclonal NBT-84868-MSM8-P0**

Artikelname	TIM3 / HAVCR2 / CD366 (Effector T-Cell Marker) Antibody, IgG2c, Clone: [TIM3/4028], Mouse, Monoclonal
Artikelnummer	NBT-84868-MSM8-P0
Hersteller Artikelnummer	84868-MSM8-P0
Alternativnummer	NBT-84868-MSM8-P0-20,NBT-84868-MSM8-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary)
Produktbeschreibung	TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cel tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 a...
Klonalität	Monoclonal
Klon-Bezeichnung	[TIM3/4028]
Molekulargewicht	60kDa

Isotyp	IgG2c
NCBI	<a href="#">84868</a>
UniProt	<a href="#">Q8TDQ0</a>
Formulierung	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 &degC followed by cooling at RT for 2