

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG2a, Clone: [UMCD2], Mouse, Monoclonal NBT-914-MSM2-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG2a, Clone: [UMCD2], Mouse, Monoclonal  |
| Artikelnummer            | NBT-914-MSM2-P0   |
| Hersteller Artikelnummer | 914-MSM2-P0   |
| Alternativnummer         | NBT-914-MSM2-P0-20,NBT-914-MSM2-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, Functional Studies, IF  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human Thymocytes/Sezary T cells   |
| Produktbeschreibung      | CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [UMCD2]   |
| Molekulargewicht         | ~50kDa  |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG2a   |
| NCBI                   | <a href="#">914</a>   |
| UniProt                | <a href="#">P06729</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.                    |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Functional Studies (Order Ab without Azide), ,Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (0.5-1ug/ml), ,Optimal dilution for a specific application should be determined. |