

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD14 (Monocyte / Macrophage Marker) Antibody, Clone: [LPSR/4180R], Rabbit, Monoclonal NBT-929-RBM14-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant CD14 (Monocyte / Macrophage Marker) Antibody, Clone: [LPSR/4180R], Rabbit, Monoclonal |
| Artikelnummer | NBT-929-RBM14-P0 |
| Hersteller Artikelnummer | 929-RBM14-P0 |
| Alternativnummer | NBT-929-RBM14-P0-20,NBT-929-RBM14-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Synthetic peptide corresponding to residues within aa1-100 of CD14 was used as an immunogen. |
| Produktbeschreibung | CD14 is receptor for endotoxin and apoptotic cells. It is expressed by monocytes/histiocytes, Langerhans cells, follicular dendritic reticulum cells, neutrophils (weakly), and endothelial cells of the liver sinusoids.CD14 is detected in myeloid leuke... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [LPSR/4180R] |
| Molekulargewicht | 40kDa |

| | |
|------------------------|--|
| NCBI | 929 |
| UniProt | P08571 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution f |