

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant CD22 / BL-CAM (B-Cell Marker) Antibody, Clone: [BLCAM/8113R], Rabbit, Monoclonal NBT-933-RBM10-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant CD22 / BL-CAM (B-Cell Marker) Antibody, Clone: [BLCAM/8113R], Rabbit, Monoclonal  |
| Artikelnummer            | NBT-933-RBM10-P0  |
| Hersteller Artikelnummer | 933-RBM10-P0  |
| Alternativnummer         | NBT-933-RBM10-P0-20,NBT-933-RBM10-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment (around aa1-200) of human BL-CAM/CD22 protein (exact sequence is proprietary)  |
| Produktbeschreibung      | Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytopla... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [BLCAM/8113R]   |
| Molekulargewicht         | 130-140kDa  |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">933</a>   |
| UniProt                | <a href="#">P20273</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution f |