

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant MUC16 / CA125 (Ovarian Carcinoma Marker) Antibody, Clone: [OCA125/2349R], Rabbit, Monoclonal NBT-94025-RBM3-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant MUC16 / CA125 (Ovarian Carcinoma Marker) Antibody, Clone: [OCA125/2349R], Rabbit, Monoclonal |
| Artikelnummer | NBT-94025-RBM3-P0 |
| Hersteller Artikelnummer | 94025-RBM3-P0 |
| Alternativnummer | NBT-94025-RBM3-P0-20,NBT-94025-RBM3-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Purified human MUC16 protein |
| Produktbeschreibung | The mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats (VNTRs). Membrane-associated and secretory Mucins are high molecular weight glycoproteins of the glycocalyx (po... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [OCA125/2349R] |
| Molekulargewicht | >2 000kDa |
| NCBI | 94025 |

| | |
|------------------------|---|
| UniProt | Q8WXI7 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |