

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant MUC16 / CA125 (Ovarian Carcinoma Marker) Antibody, Clone: [OCA125/4505R], Rabbit, Monoclonal NBT-94025-RBM5-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant MUC16 / CA125 (Ovarian Carcinoma Marker) Antibody, Clone: [OCA125/4505R], Rabbit, Monoclonal |
| Artikelnummer | NBT-94025-RBM5-P1ABX |
| Hersteller Artikelnummer | 94025-RBM5-P1ABX |
| Alternativnummer | NBT-94025-RBM5-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Synthetic peptide corresponding to CA125 residues within aa12201-12300 of CA125 was used as an immunogen. |
| Produktbeschreibung | CA125 also known as mucin 16, is a member of the mucin family glycoproteins. CA125 is a component of the ocular surface (including the cornea and conjunctiva), the respiratory tract and the female reproductive tract epithelia. Since CA125 is highly g... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [OCA125/4505R] |
| Molekulargewicht | >200kDa |

| | |
|------------------------|--|
| NCBI | 94025 |
| UniProt | Q8WXI7 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |