

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD48 (Pan Leukocyte Marker) Antibody, Clone: [CD48/8360R], Rabbit, Monoclonal NBT-962-RBM10-P1ABX

| | |
|--------------------------|--|
| Artikelname | Recombinant CD48 (Pan Leukocyte Marker) Antibody, Clone: [CD48/8360R], Rabbit, Monoclonal |
| Artikelnummer | NBT-962-RBM10-P1ABX |
| Hersteller Artikelnummer | 962-RBM10-P1ABX |
| Alternativnummer | NBT-962-RBM10-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 59-178) of human CD48 protein (exact sequence is proprietary) |
| Produktbeschreibung | Reacts with human CD48, a 45kDa glycosyl phosphatidyl-inositol (GPI)- anchored cell surface protein. CD48 is strongly expressed on lymphocytes and monocytes and weakly on granulocytes but is absent on platelets, fibroblasts, epithelium and endothelium.... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CD48/8360R] |
| Molekulargewicht | 45kDa |

| | |
|------------------------|--|
| NCBI | 962 |
| UniProt | P09326 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |