

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cdk1 / p34cdc2 Serine-Threonine Kinase Antibody, IgG2a, Clone: [A17.1.1], Mouse, Monoclonal NBT-983-MSM3-P0

| | |
|--------------------------|---|
| Artikelname | Cdk1 / p34cdc2 Serine-Threonine Kinase Antibody, IgG2a, Clone: [A17.1.1], Mouse, Monoclonal |
| Artikelnummer | NBT-983-MSM3-P0 |
| Hersteller Artikelnummer | 983-MSM3-P0 |
| Alternativnummer | NBT-983-MSM3-P0-20,NBT-983-MSM3-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Gallus, Guinea pig, Human, Mouse, Rat, Rodent, Xenopus |
| Immunogen | C-Terminal 2/3rds of Xenopus cdc2 expressed in E.coli |
| Produktbeschreibung | Recognizes a 34kDa protein (cdk1), identified as p34cdc2 (a catalytic subunit of Maturation Promoting Factor). Its epitope maps near the C-terminus of the protein and its core is thought to be LGTPNNEV (aa220-227 in murine cdc2). It shows no cross re... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [A17.1.1] |
| Molekulargewicht | 34kDa |
| Isotyp | IgG2a |

| | |
|------------------------|---|
| NCBI | 983 |
| UniProt | P06493 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Inhibits activation of p34cdc2 kinase by cyclins,,Immunofluorescence (1-2ug/ml), Immunoprecipitation, Kinase Assay, Western Blot, ,Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating |