

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cdk1 / p34cdc2 Serine-Threonine Kinase Antibody, IgG2a, Clone: [A17.1.1], Mouse, Monoclonal NBT-983-MSM3-P0**

Artikelname	Cdk1 / p34cdc2 Serine-Threonine Kinase Antibody, IgG2a, Clone: [A17.1.1], Mouse, Monoclonal
Artikelnummer	NBT-983-MSM3-P0
Hersteller Artikelnummer	983-MSM3-P0
Alternativnummer	NBT-983-MSM3-P0-20,NBT-983-MSM3-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Gallus, Guinea pig, Human, Mouse, Rat, Rodent, Xenopus
Immunogen	C-Terminal 2/3rds of Xenopus cdc2 expressed in E.coli
Produktbeschreibung	Recognizes a 34kDa protein (cdk1), identified as p34cdc2 (a catalytic subunit of Maturation Promoting Factor). Its epitope maps near the C-terminus of the protein and its core is thought to be LGTPNNEV (aa220-227 in murine cdc2). It shows no cross re...
Klonalität	Monoclonal
Klon-Bezeichnung	[A17.1.1]
Molekulargewicht	34kDa
Isotyp	IgG2a

NCBI	<a href="#">983</a>
UniProt	<a href="#">P06493</a>
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Inhibits activation of p34cdc2 kinase by cyclins,,Immunofluorescence (1-2ug/ml), Immunoprecipitation, Kinase Assay, Western Blot, ,Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating