

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, pan (Epithelial Marker) Antibody, IgG1, Clone: [KRTL/1077 + KRTH/1076], Mouse, Monoclonal NBT-MSM10-1100-P1ABX

Artikelname	Cytokeratin, pan (Epithelial Marker) Antibody, IgG1, Clone: [KRTL/1077 + KRTH/1076], Mouse, Monoclonal
Artikelnummer	NBT-MSM10-1100-P1ABX
Hersteller Artikelnummer	MSM10-1100-P1ABX
Alternativnummer	NBT-MSM10-1100-P1ABX-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human KRT77 and KRT76 protein
Produktbeschreibung	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which include CK1, CK3, CK4, CK5, CK6, CK8...
Klonalität	Monoclonal
Klon-Bezeichnung	[KRTL/1077 + KRTH/1076]
Molekulargewicht	67kDa (CK1), 64kDa (CK3), 59kDa (CK4), 58kDa (CK5), 56kDa (CK6), 52kDa (CK8), 56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19)

Isotyp	IgG1
NCBI	374454
UniProt	Q7Z794
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f