

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myeloid Cell Marker (Macrophage / Granulocyte Marker) Antibody, IgG1, Clone: [BM-1], Mouse, Monoclonal NBT-MSM1-61-P1ABX

| | |
|--------------------------|---|
| Artikelname | Myeloid Cell Marker (Macrophage / Granulocyte Marker) Antibody, IgG1, Clone: [BM-1], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM1-61-P1ABX |
| Hersteller Artikelnummer | MSM1-61-P1ABX |
| Alternativnummer | NBT-MSM1-61-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human peripheral blood mononuclear cells |
| Produktbeschreibung | Recognizes 183kDa protein with DNA-binding characteristics, which is identified as a myeloid specific antigen. It reacts with myeloid precursor cells and granulocytes in bone marrow. Its antigen appears to be restricted to M2 and M3 acute myelogenous... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BM-1] |
| Molekulargewicht | 183kDa |
| Isotyp | IgG1 |

| | |
|------------------------|--|
| NCBI | Not Known |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (No special pretreatment is required for staining of formalin/paraffin tissues.)Optimal dilution for a specific app |