

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Myeloid Cell Marker (Macrophage / Granulocyte Marker) Antibody, IgG1, Clone: [BM-1], Mouse, Monoclonal NBT-MSM1-61-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Myeloid Cell Marker (Macrophage / Granulocyte Marker) Antibody, IgG1, Clone: [BM-1], Mouse, Monoclonal  |
| Artikelnummer            | NBT-MSM1-61-P1ABX   |
| Hersteller Artikelnummer | MSM1-61-P1ABX   |
| Alternativnummer         | NBT-MSM1-61-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human peripheral blood mononuclear cells  |
| Produktbeschreibung      | Recognizes 183kDa protein with DNA-binding characteristics, which is identified as a myeloid specific antigen. It reacts with myeloid precursor cells and granulocytes in bone marrow. Its antigen appears to be restricted to M2 and M3 acute myelogenous... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [BM-1]  |
| Molekulargewicht         | 183kDa  |
| Isotyp                   | IgG1  |

|                        |  |
|------------------------|--|
| NCBI                   | Not Known  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (No special pretreatment is required for staining of formalin/paraffin tissues.)Optimal dilution for a specific app |