

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CA19-9/Sialyl Lewisa (GI Tumor Marker) Antibody, IgM, Clone: [SPM588], Mouse, Monoclonal NBT-MSM1X-25-P0

| | |
|--------------------------|---|
| Artikelname | CA19-9/Sialyl Lewisa (GI Tumor Marker) Antibody, IgM, Clone: [SPM588], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM1X-25-P0 |
| Hersteller Artikelnummer | MSM1X-25-P0 |
| Alternativnummer | NBT-MSM1X-25-P0-20,NBT-MSM1X-25-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Precipitin lines obtained after immuno-diffusion using MAb 116-NS-19-9 and mucins isolated from an ovarian cyst of a Lewis A+B-patient (OLe). |
| Produktbeschreibung | CA19-9, a carbohydrate epitope expressed on a high MW (400kDa) mucin glycoprotein, is a sialyl Lewisa structure which is synthesized from type 1 blood group precursor chains and is present in individuals expressing the Lewis a and/or Lewis b blood grou... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM588] |
| Molekulargewicht | >400kDa |

| | |
|------------------------|--|
| Isotyp | IgM |
| NCBI | Not Known |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(No special pretreatment is required for the immunohistochemical staining of formalin-fixed, paraffin-embedded tissu |