

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **HSV1 (Herpes Simplex Virus Type I) Antibody, IgG1, Clone: [10A3], Mouse, Monoclonal NBT-MSM2-1793-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | HSV1 (Herpes Simplex Virus Type I) Antibody, IgG1, Clone: [10A3], Mouse, Monoclonal   |
| Artikelnummer            | NBT-MSM2-1793-P0  |
| Hersteller Artikelnummer | MSM2-1793-P0  |
| Alternativnummer         | NBT-MSM2-1793-P0-20,NBT-MSM2-1793-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Virus   |
| Immunogen                | Detergent-solubilized herpes simplex virus (HSV) type 1 infected cells  |
| Produktbeschreibung      | The antibody reacts with HSV type 1 specific antigen. It is suitable for detection of HSV in human cellular material obtained from superficial lesions or biopsies and for the early identification of HSV in infected tissue cultures. The herpes simplex... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [10A3]  |
| Isotyp                   | IgG1  |

|                        |   |
|------------------------|---|
| NCBI                   | Not Applicable  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution f |