

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Phosphotyrosine (P-Tyr) Antibody, IgG2b, Clone: [PY793], Mouse, Monoclonal NBT-MSM2-793-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Phosphotyrosine (P-Tyr) Antibody, IgG2b, Clone: [PY793], Mouse, Monoclonal  |
| Artikelnummer            | NBT-MSM2-793-P1ABX  |
| Hersteller Artikelnummer | MSM2-793-P1ABX  |
| Alternativnummer         | NBT-MSM2-793-P1ABX-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC   |
| Spezies Reaktivität      | All   |
| Immunogen                | Phosphotyrosine conjugated to BSA   |
| Produktbeschreibung      | Protein phosphorylation is a fundamental event in the regulation of a large number of intracellular processes. Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [PY793]   |
| Molekulargewicht         | Depends upon the phosphorylated target  |
| Isotyp                   | IgG2b   |

|                        |  |
|------------------------|--|
| NCBI                   | Not Applicable   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (No pretreatment is required for staining of formalin-fixed, paraffin-embedded tissues)Optimal dilution for a speci |