

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Lewis Y (Tumor Marker) Antibody, IgG1, Clone: [LWY/1463], Mouse, Monoclonal NBT-MSM3-1463-P1ABX

| | |
|--------------------------|---|
| Artikelname | Lewis Y (Tumor Marker) Antibody, IgG1, Clone: [LWY/1463], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM3-1463-P1ABX |
| Hersteller Artikelnummer | MSM3-1463-P1ABX |
| Alternativnummer | NBT-MSM3-1463-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human colon carcinoma cells |
| Produktbeschreibung | This antibody recognizes a carbohydrate epitope present on tumor-associated Lewis Y antigen (Fucalpha1-2Galbeta1-4/3[Fucalpha1-3/4]GlcNAcbeta-). Lewis Y is expressed in large bowel tumors and colorectal carcinomas. It may be useful in the classificat... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [LWY/1463] |
| Molekulargewicht | Multiple |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| NCBI | Not Applicable |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |