

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Epstein-Barr Virus (LMP-1) Antibody, IgG1, Clone: [CS2], Mouse, Monoclonal NBT-MSM3-1742-P1ABX

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Epstein-Barr Virus (LMP-1) Antibody, IgG1, Clone: [CS2], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM3-1742-P1ABX |
| Hersteller Artikelnummer | MSM3-1742-P1ABX |
| Alternativnummer | NBT-MSM3-1742-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, WB |
| Spezies Reaktivität | Virus |
| Immunogen | Recombinant fusion protein containing the sequence of bacterial beta-galactosidase and the carboxyl half of EBV-encoded LMP |
| Produktbeschreibung | This antibody is a mixture of four different monoclonal antibodies. This antibody is specific to 60kDa latent membrane protein (LMP-1) encoded by the BNLF1 gene of the EBV. Each clone reacts with different epitopes on the hydrophilic C-terminus of th... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CS2] |
| Molekulargewicht | 60kDa |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG1 |
| NCBI | Not Applicable |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunofluorescence (0.5-1.0ug/ml), Western Blotting (0.5-1.0ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min |