

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Cytokeratin 5/6/18 Antibody, IgG1, Clone: [LP34], Mouse, Monoclonal NBT-MSM36-2036-P0

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 5/6/18 Antibody, IgG1, Clone: [LP34], Mouse, Monoclonal   |
| Artikelnummer            | NBT-MSM36-2036-P0   |
| Hersteller Artikelnummer | MSM36-2036-P0   |
| Alternativnummer         | NBT-MSM36-2036-P0-20,NBT-MSM36-2036-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Bovine, Human, Rat  |
| Immunogen                | Detergent-insoluble fraction of psoriatic human epidermis   |
| Produktbeschreibung      | It recognizes polypeptides of 58kDa, 56kDa, and 45kDa, identified as cytokeratin 5, 6, and 18 respectively. It shows no reaction with keratin 1, 8, or 19. This MAb shows a broad pattern of reactivity with epithelial tissues, from simple glandular epi... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [LP34]  |
| Molekulargewicht         | 45.58kDa  |
| Isotyp                   | IgG1  |
| NCBI                     | Multiple  |

|                        |   |
|------------------------|---|
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |