

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HepPar-1 (Hepatocellular Marker) Antibody, IgG1, Clone: [HepPar1], Mouse, Monoclonal NBT-MSM4-966-P0

Artikelname	HepPar-1 (Hepatocellular Marker) Antibody, IgG1, Clone: [HepPar1], Mouse, Monoclonal
Artikelnummer	NBT-MSM4-966-P0
Hersteller Artikelnummer	MSM4-966-P0
Alternativnummer	NBT-MSM4-966-P0-20,NBT-MSM4-966-P0-100
Hersteller	NeoBiotechnologies
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC
Spezies Reaktivität	Canine, Human
Immunogen	Extract of a formalin-fixed, rejected-allograft of a human liver
Produktbeschreibung	Hepatocyte Paraffin 1 or HepPar1 localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio-carcinomas. HCCs occur primarily in the s...
Klonalität	Monoclonal
Klon-Bezeichnung	[HepPar1]
Molekulargewicht	Not Known
Isotyp	IgG1

NCBI	Not Known
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95°C followed by cooling at RT for 20