

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant HPV-16 (Human Papilloma Virus 16) Antibody, IgG1, Clone: [rHPV16L1/1058], Mouse, Monoclonal NBT-MSM7-2057-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant HPV-16 (Human Papilloma Virus 16) Antibody, IgG1, Clone: [rHPV16L1/1058], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM7-2057-P0 |
| Hersteller Artikelnummer | MSM7-2057-P0 |
| Alternativnummer | NBT-MSM7-2057-P0-20,NBT-MSM7-2057-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Virus |
| Immunogen | Recombinant full-length protein corresponding to Human papillomavirus HPV16 L1. |
| Produktbeschreibung | Reacts with a protein of 57kDa, identified as the L1 protein of human papilloma virus type 16 (HPV-16). It is the major capsid protein of HPV-16. Infection with specific types of HPV has been associated with an increased risk of developing cervical n... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [rHPV16L1/1058] |
| Molekulargewicht | 57kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | Not Applicable |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistology (Formalin-paraffin) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specif |