

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) Antibody, Clone: [KRTL/1577R], Rabbit, Monoclonal NBT-RBM21-1577-P1ABX

| | |
|--------------------------|---|
| Artikelname | Recombinant Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) Antibody, Clone: [KRTL/1577R], Rabbit, Monoclonal |
| Artikelnummer | NBT-RBM21-1577-P1ABX |
| Hersteller Artikelnummer | RBM21-1577-P1ABX |
| Alternativnummer | NBT-RBM21-1577-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Canine, Human |
| Immunogen | Recombinant human KRT77 protein fragment |
| Produktbeschreibung | This MAb recognizes the 56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamil... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRTL/1577R] |
| Molekulargewicht | 56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19) |

| | |
|------------------------|---|
| NCBI | 374454 |
| UniProt | Q7Z794 |
| Formulierung | 200ug/ml of Ab Purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |