

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Cytokeratin, Basic (Type II or HMW) Antibody, Clone: [KRTH/2147R], Rabbit, Monoclonal NBT-RBM31-2147-P0

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Recombinant Cytokeratin, Basic (Type II or HMW) Antibody, Clone: [KRTH/2147R], Rabbit, Monoclonal |
| Artikelnummer | NBT-RBM31-2147-P0 |
| Hersteller Artikelnummer | RBM31-2147-P0 |
| Alternativnummer | NBT-RBM31-2147-P0-20,NBT-RBM31-2147-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Solubilized keratin extract from human stratum corneum |
| Produktbeschreibung | This antibody recognizes CK1, CK5, CK10 and CK14. In normal epithelia, it stains stratified epithelia, myoepithelial cells and basal cells in the prostate gland and bronchi. This MAb shows no reactivity with hepatocytes, pancreatic acinar cells, prox... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRTH/2147R] |
| Molekulargewicht | 50-67kDa |
| NCBI | 51350 |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | Q01546 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2 ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1m |