

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Cytokeratin, Multi (Epithelial Marker) Antibody, Clone: [KRT/1877R], Rabbit, Monoclonal NBT-RBM37-1877-P0

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Recombinant Cytokeratin, Multi (Epithelial Marker) Antibody, Clone: [KRT/1877R], Rabbit, Monoclonal |
| Artikelnummer | NBT-RBM37-1877-P0 |
| Hersteller Artikelnummer | RBM37-1877-P0 |
| Alternativnummer | NBT-RBM37-1877-P0-20,NBT-RBM37-1877-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Bovine, Frog, Goat, Guinea pig, Human, Monkey, Mouse, Porcine, Rat |
| Immunogen | Keratin-enriched preparation from cultured human epithelial cells |
| Produktbeschreibung | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, including 59kDa (CK4), 58kDa (CK5), 56kDa (CK6), 52... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT/1877R] |
| Molekulargewicht | Multiple |
| NCBI | 3851 |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P02538 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA |