

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Ep-CAM / CD326 (Rat) Antibody, IgG1, Clone: [Epcam/1158], Mouse, Monoclonal NBT-RT171577-MSM1-BC0

|                          |   |
|--------------------------|---|
| Artikelname              | Ep-CAM / CD326 (Rat) Antibody, IgG1, Clone: [Epcam/1158], Mouse, Monoclonal   |
| Artikelnummer            | NBT-RT171577-MSM1-BC0   |
| Hersteller Artikelnummer | RT171577-MSM1-BC0   |
| Alternativnummer         | NBT-RT171577-MSM1-BC0-0.1,NBT-RT171577-MSM1-BC0-0.5,NBT-RT171577-MSM1-BC0-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length Rat Epcam protein   |
| Produktbeschreibung      | Epitopes of Epcam/1158 and Epcam/1159 monoclonal antibodies are different. EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expre... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [Epcam/1158]  |
| Molekulargewicht         | 40-43kDa  |

|                        |  |
|------------------------|--|
| Isotyp                 | IgG1   |
| NCBI                   | <a href="#">171577</a>   |
| UniProt                | <a href="#">O55159</a>   |
| Formulierung           | Bioreactor Concentrate with 0.05% Azide. Contact us if you require it in a different format.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1:50-1:100, 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |