

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Rat) Antibody, IgG1, Clone: [EpCAM/1159], Mouse, Monoclonal NBT-RT171577-MSM2-BC0

| | |
|--------------------------|---|
| Artikelname | Ep-CAM / CD326 (Rat) Antibody, IgG1, Clone: [EpCAM/1159], Mouse, Monoclonal |
| Artikelnummer | NBT-RT171577-MSM2-BC0 |
| Hersteller Artikelnummer | RT171577-MSM2-BC0 |
| Alternativnummer | NBT-RT171577-MSM2-BC0-0.1,NBT-RT171577-MSM2-BC0-0.5,NBT-RT171577-MSM2-BC0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length Rat Epcam protein |
| Produktbeschreibung | Epitopes of Epcam/1158 and Epcam/1159 monoclonal antibodies are different. EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expre... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [EpCAM/1159] |
| Molekulargewicht | 40-43kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 171577 |
| UniProt | O55159 |
| Formulierung | Bioreactor Concentrate with 0.05% Azide. Contact us if you require it in a different format. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1:50-1:100, 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |