

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### FTO Antibody, Unconjugated, Rabbit, Polyclonal ROC-600-401-BH8

|                          |   |
|--------------------------|---|
| Artikelname              | FTO Antibody, Unconjugated, Rabbit, Polyclonal  |
| Artikelnummer            | ROC-600-401-BH8   |
| Hersteller Artikelnummer | 600-401-BH8   |
| Alternativnummer         | ROC-600-401-BH8   |
| Hersteller               | Rockland Immunochemicals  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, IF, IHC, WB  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Anti-FTO antibody was prepared from whole rabbit serum produced by repeated immunizations with a 15 amino acid synthetic peptide from near the N-terminus of human FTO. |
| Konjugation              | Unconjugated  |
| Klonalität               | Polyclonal  |
| Konzentration            | 1 mg/mL by UV absorbance at 280 nm  |
| NCBI                     | <a href="#">79068</a>   |
| UniProt                  | <a href="#">Q9C0B1</a>  |
| Puffer                   | 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2   |
| Formulierung             | Liquid (sterile filtered)   |

|                        |   |
|------------------------|---|
| Target-Kategorie       | Human   |
| Antibody Type          | Primary Antibody  |
| Application Verdünnung | ELISA: 1:10,000 - 1:20,000, IHC: 2.5 µg/mL, IF Microscopy: 20 µg/mL, WB: 1-2 µg/mL  |
| Anwendungsbeschreibung | Anti-FTO Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 58 kDa in Western Blots of specific |