

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### NOTUM Antibody, Unconjugated, Rabbit, Polyclonal ROC-600-401-DD7

|                          |  |
|--------------------------|--|
| Artikelname              | NOTUM Antibody, Unconjugated, Rabbit, Polyclonal   |
| Artikelnummer            | ROC-600-401-DD7  |
| Hersteller Artikelnummer | 600-401-DD7  |
| Alternativnummer         | ROC-600-401-DD7  |
| Hersteller               | Rockland Immunochemicals   |
| Wirt                     | Rabbit   |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, IF, IHC, WB   |
| Spezies Reaktivität      | Human, Mouse, Rat  |
| Immunogen                | Anti-NOTUM antibody was prepared from whole rabbit serum produced by repeated immunizations with a 19 amino acid synthetic peptide near the N-terminus of human NOTUM. |
| Konjugation              | Unconjugated   |
| Klonalität               | Polyclonal   |
| Konzentration            | 1 mg/mL by UV absorbance at 280 nm   |
| NCBI                     | <a href="#">147111</a>   |
| UniProt                  | <a href="#">Q6P988</a>   |
| Puffer                   | 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2  |
| Formulierung             | Liquid (sterile filtered)  |

|                        |  |
|------------------------|--|
| Target-Kategorie       | Human  |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | ELISA: 1:5,000 - 1:10,000, IHC: 2.5 µg/mL, IF Microscopy: 20 µg/mL, WB: 0.5-1 µg/mL  |
| Anwendungsbeschreibung | Anti-NOTUM Antibody has been tested for use in ELISA, Western Blotting, Immunohistochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 56 kDa in Western Blots of specifi |