

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Angiostatin K1-3/PLG Antibody, Rabbit, Polyclonal BOB-RP1027

Artikelname	Anti-Angiostatin K1-3/PLG Antibody, Rabbit, Polyclonal
Artikelnummer	BOB-RP1027
Hersteller Artikelnummer	RP1027
Alternativnummer	BOB-RP1027-10UG,BOB-RP1027-100UG
Hersteller	Boster Bio
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Immunogen	E. coli-derived human Angiostatin K1-3 recombinant protein (Position: C103-C352).
Produktbeschreibung	Boster Bio Anti-Angiostatin K1-3/PLG Antibody catalog RP1027. Tested in ELISA, IHC, WB applications. This antibody reacts with Human....
Klonalität	Polyclonal
Konzentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Molekulargewicht	90569 MW
UniProt	P00747

Puffer	Each vial contains 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify Carrier Free or No BSA in your order note.
Reinheit	Immunogen affinity purified.
Formulierung	Lyophilized
Application Verdünnung	Immunohistochemistry (Paraffin-embedded Section), 0.5-1µg/ml, Human, By Heat ELISA , 0.1-0.5µg/ml, Human, -Western blot, 0.1-0.5µg/ml, Human
Anwendungsbeschreibung	WB: The detection limit for PLG is approximately 0.5ng/lane under reducing conditions. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have no