

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-CD3 epsilon (activation epitope) Purified Monoclonal Antibody, Clone: [Clone: APA1/1], Mouse BOB-M02675-4

| | |
|--------------------------|---|
| Artikelname | Anti-CD3 epsilon (activation epitope) Purified Monoclonal Antibody, Clone: [Clone: APA1/1], Mouse |
| Artikelnummer | BOB-M02675-4 |
| Hersteller Artikelnummer | M02675-4 |
| Alternativnummer | BOB-M02675-4-0.1MG |
| Hersteller | Boster Bio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, ICC, IHC-Fr, IP, WB |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Purified human CD3 proteins isolated from thymus. The mouse monoclonal antibody APA1/1 recognizes an activation-dependent intracellular epitope of CD3 epsilon. Exposure of the epitope precedes CD3 phosphorylation and recruitment and activation of ZAP70, |
| Produktbeschreibung | Boster Bio Anti-CD3 epsilon (activation epitope) Purified Monoclonal Antibody (Catalog M02675-4). Tested in Flow Cytometry, IP, WB, IHC-F, ICC application(s). This antibody reacts with Mouse, Human.... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/ml |

| | |
|------------------------|--|
| Klon-Bezeichnung | [Clone: APA1/1] |
| Molekulargewicht | 23147 MW |
| UniProt | P07766 |
| Puffer | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Reinheit | Purified by protein-A affinity chromatography. |
| Formulierung | Liquid |
| Application Verdünnung | Flow cytometry: Intracellular staining, recommended dilution: 1-4 µg/ml, positive control: human T cells stimulated with anti-CD3 (MEM-57) antibody (1 µg/ml). Sample preparation: At the end of stimulation of T cells, perform staining of surface markers (|