

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10], IgG1, Clone: [Clone: Clone: 4E10], Mouse BOB-M03295

|                          |  |
|--------------------------|--|
| Artikelname              | Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10], IgG1, Clone: [Clone: Clone: 4E10], Mouse  |
| Artikelnummer            | BOB-M03295   |
| Hersteller Artikelnummer | M03295   |
| Alternativnummer         | BOB-M03295-0.1MG   |
| Hersteller               | Boster Bio   |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, FC, ICC, IF, IHC-P, WB  |
| Spezies Reaktivität      | Human  |
| Immunogen                | PD-L2 antibody was raised against the extracellular domain of human PD-L2.   |
| Produktbeschreibung      | Boster Bio Anti-PDL2 PDCD1LG2 Monoclonal Antibody [4E10] (Catalog M03295). Tested in ELISA, WB, IHC-P, ICC, IF, Flow Cytometry applications. This antibody reacts with Human.... |
| Klonalität               | Monoclonal   |
| Konzentration            | 1 mg/mL  |
| Klon-Bezeichnung         | [Clone: Clone: 4E10]   |
| Isotyp                   | IgG1   |

|                        |   |
|------------------------|---|
| UniProt                | <a href="#">P9BQ51</a>  |
| Puffer                 | PD-L2 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol.   |
| Reinheit               | PD-L2 Antibody is supplied as protein A purified IgG1.  |
| Formulierung           | Liquid  |
| Application Verdünnung | PD-L2 antibody can be used for detection of PD-L2 by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples, Im |