

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Angiostatin K1-3/PLG Antibody Picoband, Rabbit, Polyclonal BOB-RP1027

| | |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Anti-Angiostatin K1-3/PLG Antibody Picoband, Rabbit, Polyclonal |
| Artikelnummer | BOB-RP1027 |
| Hersteller Artikelnummer | RP1027 |
| Alternativnummer | BOB-RP1027-100UG |
| Hersteller | Boster Bio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | E. coli-derived human Angiostatin K1-3 recombinant protein (Position: C103-C352). |
| Produktbeschreibung | Boster Bio Anti-Angiostatin K1-3/PLG Antibody catalog RP1027. Tested in ELISA, IHC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong... |
| Klonalität | Polyclonal |
| Konzentration | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml. |
| Molekulargewicht | 95 kDa |
| UniProt | P00747 |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer | Each vial contains 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify Carrier Free or No BSA in your order note. |
| Reinheit | Immunogen affinity purified. |
| Formulierung | Lyophilized |
| Application Verdünnung | Immunohistochemistry (Paraffin-embedded Section), 0.5-1µg/ml, Human, By Heat ELISA , 0.1-0.5µg/ml, Human, -Western blot, 0.1-0.5µg/ml, Human |