

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cy3-conjugated AffiniPure(TM) Fab Fragment Bovine Anti-Goat IgG, Fc Fragment Specific JIM-805-167-008**

|                          |   |
|--------------------------|---|
| Artikelname              | Cy3-conjugated AffiniPure(TM) Fab Fragment Bovine Anti-Goat IgG, Fc Fragment Specific   |
| Artikelnummer            | JIM-805-167-008   |
| Hersteller Artikelnummer | 805-167-008   |
| Alternativnummer         | JIM-805-167-008   |
| Hersteller               | Jackson ImmunoResearch  |
| Wirt                     | Bovine  |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Goat  |
| Konjugation              | Cyanine 3   |
| Klonalität               | Polyclonal  |
| Puffer                   | Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free). Preservative: 0.05% Sodium Azide   |
| Reinheit                 | AffiniPure  |
| Formulierung             | Freeze-dried solid  |
| Lagerung                 | Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with the indicated volume of dH2O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended... |

|                        |   |
|------------------------|---|
| Antibody Type          | Secondary Antibody  |
| Application Verdünnung | _x000D_ 1:100 - 1:800 for sequential labeling applications._x000D_<br>_x000D_ To complex with primary antibody in solution, use<br>1:1 weight ratio of Fab:primary antibody (3:1 molar ratio). Vortex<br>and incubate for 30 minutes at room temperature... |