

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Rhodamine Red-X-conjugated AffiniPure(TM) Bovine Anti-Goat IgG (H+L) (min X Bov,Ck,GP,Sy Hms,Hrs,Hu,Ms,Rb,Rat Sr Prot) JIM-805-295-180**

|                          |   |
|--------------------------|---|
| Artikelname              | Rhodamine Red-X-conjugated AffiniPure(TM) Bovine Anti-Goat IgG (H+L) (min X Bov,Ck,GP,Sy Hms,Hrs,Hu,Ms,Rb,Rat Sr Prot)  |
| Artikelnummer            | JIM-805-295-180   |
| Hersteller Artikelnummer | 805-295-180   |
| Alternativnummer         | JIM-805-295-180   |
| Hersteller               | Jackson ImmunoResearch  |
| Wirt                     | Bovine  |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Goat  |
| Konjugation              | Rhodamine Red™ -X   |
| Klonalität               | Polyclonal  |
| Puffer                   | Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free). Preservative: 0.05% Sodium Azide   |
| Reinheit                 | AffiniPure  |
| Formulierung             | Freeze-dried solid  |
| Lagerung                 | Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with the indicated volume of dH2O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended... |

|                        |   |
|------------------------|---|
| Antibody Type          | Secondary Antibody  |
| Application Verdünnung | 1:50 - 1:200 for most applications<br>Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be... |