

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Sheep anti Dog IgG (Fc specific), Clone: [Polyclonal], Monoclonal NMB-SHAD/IGG(FC)**

|                          |  |
|--------------------------|--|
| Artikelname              | Sheep anti Dog IgG (Fc specific), Clone: [Polyclonal], Monoclonal  |
| Artikelnummer            | NMB-SHAD/IGG(FC)   |
| Hersteller Artikelnummer | ShAD/IgG(Fc)   |
| Alternativnummer         | NMB-SHAD/IGG(FC)   |
| Hersteller               | NordicMubio  |
| Wirt                     | Sheep  |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, ICC   |
| Spezies Reaktivität      | Canine   |
| Produktbeschreibung      | The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunolectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal dog serum a single precipitin line is obtained which shows a ...   |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [Polyclonal]   |
| Puffer                   | Delipidated, heat inactivated, lyophilized, stable whole antiserum. 0.01% sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added. Reconstitute the lyophilized an |

|                        |  |
|------------------------|--|
| Quelle                 | Highly purified normal IgG isolated from pooled dog serum was used for immunization of the sheep. Freunds complete adjuvant is used in the first step of the immunization procedure.   |
| Formel                 | Delipidated, heat inactivated, lyophilized, stable whole antiserum. 0.01% sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added.                                 |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat) |